



# Mec1-Dependent Phosphorylation of the Scc3 Subunit of Cohesin during Mitosis in Budding Yeast

Nathalie Grandin, Michel Charbonneau

## ► To cite this version:

Nathalie Grandin, Michel Charbonneau. Mec1-Dependent Phosphorylation of the Scc3 Subunit of Cohesin during Mitosis in Budding Yeast. *Advances in Bioscience and Biotechnology*, 2015, 06 (03), pp.153-163. 10.4236/abb.2015.63015 . hal-01154292

**HAL Id: hal-01154292**

**<https://univ-tours.hal.science/hal-01154292>**

Submitted on 21 May 2015

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Mec1-Dependent Phosphorylation of the Scc3 Subunit of Cohesin during Mitosis in Budding Yeast

Nathalie Grandin, Michel Charbonneau\*

Université François-Rabelais de Tours, CNRS, GICC UMR 7292, Tours, France  
Email: \*[michel.charbonneau@univ-tours.fr](mailto:michel.charbonneau@univ-tours.fr)

Received 21 February 2015; accepted 8 March 2015; published 17 March 2015

Copyright © 2015 by authors and Scientific Research Publishing Inc.  
This work is licensed under the Creative Commons Attribution International License (CC BY).  
<http://creativecommons.org/licenses/by/4.0/>



Open Access

---

## Abstract

Cohesin is an evolutionary conserved complex that controls chromosome segregation during mitosis. Here we show that, in response to DNA damage, *Saccharomyces cerevisiae* Scc3, one of the major regulatory subunits of the Smc1-Smc3-Scc1 cohesin ring, is phosphorylated on S/T-Q residues. This event depended on the Mec1 checkpoint kinase as well as on cell cycle arrest triggered by the DNA damage checkpoint network. This phosphorylation event also took place during mitosis of an unperturbed cell cycle. The present finding that *S. cerevisiae* Scc3 is phosphorylated during mitosis represents a potentially important new regulatory step in cohesin's mitotic functions.

## Keywords

Cohesin, S/T-Q Phosphorylation, Mec1 and Tel1, DNA Damage, Cell Cycle, Budding Yeast

---

## 1. Introduction

Cohesin, a tripartite ring composed of the Smc1, Smc3 and kleisin (Scc1/Mcd1 in the budding yeast *Saccharomyces cerevisiae*, RAD21/SCC1 in humans) subunits, is essential in all eukaryotic organisms studied to date both for establishing sister chromatid cohesion during DNA replication, maintaining them together until metaphase, and allowing their timely separation during anaphase of mitosis [1]-[4]. In addition to its role in controlling cohesion between chromatids [5]-[7], cohesin plays important roles in chromosome condensation, DNA repair, transcription regulation, chromatin structure, homolog pairing, mono-orientation of sister kinetochore and maintenance of centrosome number during mitosis [8]-[10]. Early experiments in both fission and budding yeasts have identified mutants of cohesin subunits, particularly Scc1/Rad21 and Wapl1/Rad61, which are hy-

---

\*Corresponding author.

persensitive to DNA damage, probably due to defective DNA repair resulting from inability of cohesin to maintain sister chromatid cohesion [5]. More recently, it has been shown that in response to DNA damage, the Scc1 budding yeast subunit of cohesin is phosphorylated by Chk1 in order to provide *de novo* establishment of sister chromatid cohesion [11]. In vertebrates, cohesin also appears to be regulated by phosphorylation in response to DNA damage, but in this case, phosphorylation occurs on Smc1 and Smc3 and is mediated by the ATM and ATR kinases [12] [13].

In the present study, we have set out to detect possible phosphorylation of *S. cerevisiae* cohesin on S/T-Q residues, representing a consensus for phosphorylation by the MEC1 and TEL1 kinases (ATR and ATM in higher eukaryotes, respectively [14].

## 2. Materials and Methods

### 2.1. Strains and Plasmids

Yeast strains used in this study were derivatives of BF264-15Daub (*ade1 his2 leu2-3, 112 trp1-1a ura3Dns*), as described previously [15]. Yeast cultures were grown at the indicated temperatures in YEP (1% yeast extract, 2% bacto-peptone, 0.005% adenine, 0.005% uracil) supplemented with 2% glucose (YEPD), sucrose or galactose, or in selective minimal medium. All strains were made isogenic by back crossing at least five times against our genetic background. Strain origins, prior to back crossing and, in some cases, modification of the selectable genetic marker, were as follows. The *scc1::KanMX4/SCC1*, *tel1::KanMX4* and *chk1::KanMX4* strains were purchased at Euroscarf (Frankfurt, Germany). The *mec1::TRP1* and *sml1::KanMX4* strains were from the Emili laboratory [16]. The *rad24::URA3* strain was from the Friedberg laboratory [17]. The *rad53-kd* (K227A) strain was from the Heyer laboratory [18]. The *cdc13-1* strain was from the Hartwell laboratory [19]. The *mec1-kd* (D2224A, N2229K) plasmid was from the Matsuura laboratory [20]. The *mad2::URA3* disruption plasmid was from the Murray laboratory [21].

All constructs for tagging genes (unique, endogenous copy) with 2 HA in 3', 2 Myc in 5' or 3', 9 or 18 Myc in 3' were made by using Polymerase Chain Reaction (PCR) to adapt the relevant restriction sites to the sequence of the genes. Details of the constructs will be made available upon request. The Myc<sub>18</sub>-MEC1 construct was from the Matsuura laboratory [20].

### 2.2. Immunoprecipitation

Techniques for block and release experiments, cell extract preparation, immunoprecipitation and immunoblotting (analyzed using a FUJI-LAS4000) have been described previously [22]. Mouse monoclonal anti-HA raw ascites fluid, clone 16B12 (Covance), and mouse monoclonal anti-HA 12CA5 antibody (Roche Diagnostics) were used for immunoprecipitation and immunoblotting, respectively. Myc-tagged proteins were visualized after immunoprecipitation and Western blotting with monoclonal anti-Myc antibody, clone 9E10 (Roche Diagnostics). Anti-Phospho-(Ser/Thr) ATM/ATR substrate antibody was from Cell Signaling Technology (cat.#2851).

### 2.3. Mass Spectrometry Analysis

Mass spectrometry analysis was performed at the "Plate-forme d'Analyse Intégrative des Biomarqueurs" (INRA, Nouzilly, France). The proteins in polyacrylamide gel corresponding to the P-S/T-Q reactive band on the immunoblot membrane were analyzed by nanoflow liquid chromatography-tandem mass spectrometry (nanoLC-MS/MS). The gel slice was washed in water/acetonitrile (1:1) for 5 min and in acetonitrile for 10 min. Cysteine reduction and alkylation were performed by successive incubations in 10 mM dithiothreitol/50 mM NH<sub>4</sub>HCO<sub>3</sub> for 30 min at 56°C and 55 mM iodoacetamide/50 mM NH<sub>4</sub>HCO<sub>3</sub> for 20 min at room temperature in the dark. Gel slices were washed by incubation in 50 mM NH<sub>4</sub>HCO<sub>3</sub>/acetonitrile (1:1) for 10 min and by incubation in acetonitrile for 15 min. Proteins were digested overnight in 25 mM NH<sub>4</sub>HCO<sub>3</sub> with 12.5 ng/μl trypsin (Sequencing Grade, Roche, Paris). The resulting peptides were extracted from gel by successive incubations in 0.1% formic acid (FA)/acetonitrile (1:1) for 10 min and in acetonitrile for 5 min. The extracts were pooled, dried and submitted to phosphopeptide enrichment using HyperSep MicroTip titanium dioxide (TiO<sub>2</sub>), (Thermo-Fisher). Enriched peptides were analyzed by nanoflow liquid chromatography-tandem mass spectrometry (nanoLC-MS/MS). The experiments were performed on a LTQ Orbitrap Velos mass spectrometer (Thermo Fisher Scientific, Bremen, Germany) coupled to an Ultimate® 3000 RSLC Ultra High Pressure Liquid Chroma-

tographer (Dionex, Amsterdam, The Netherlands). Enriched peptides were loaded on an LCPackings trap column (Acclaim PepMap 100 C<sub>18</sub>, 100 µm inner diameter × 2 cm long, 3 µm particles, 100 Å pores). Mobile phases consisted of (A) 0.1% formic acid (FA), 97.9% water, 2% acetonitrile (v/v/v) and (B) 0.1% FA, 15.9% water, 84% acetonitrile (v/v/v). Peptides were desalted for 10 min at 5 µL/min with 4% solvent B and separated using a LCPackings nano-column (Acclaim PepMap C<sub>18</sub>, 75 µm inner diameter × 50 cm long, 3 µm particles, 100 Å pores). The gradient consisted of 4% - 55% B for 90 min. The nanoflow rate was set at 300 nL/min.


Data were acquired using Xcalibur software (version 2.1; Thermo Fisher Scientific, San Jose, CA). The instrument was operated in positive ionization mode. The Orbitrap analyzer recorded signals between 300 and 1800 *m/z* with a resolution set to 60,000 in MS. The minimum MS signal for triggering MS/MS was set to 500 counts. During MS/MS, fragmentation with Collision Induced Dissociation (CID) and detection occurred in the linear ion trap analyzer using the “Neutral Loss” and “MultiStage Activation (MSA)” MS methods. Neutral loss masses specified for fragmentations were 49 and 32.7 (for di and tricharged monophosphopeptide) and 98 and 65.3 (for di and tricharged diphosphopeptide). Precursor selection window was 2 *m/z* (1 microscan) for MS/MS. Data acquiring was automatically done with a profile MS full scan, a centroid MS/MS scan on the 5 most intense peptide ions with charge states ≥ 2 and MS<sup>3</sup> fragmentation on the three most intense detected peaks using neutral loss detection. The maximum ion accumulation times allowed were 500 ms for full scans and 25 ms for CID-MS/MS. Target ion quantity for FT full MS was 1e6 and for MS<sup>n</sup> it was 1e4. The resulting fragment ions were scanned at the “normal scan rate” with *q* = 0.25 activation and activation time of 10 ms. Dynamic exclusion was activated for 30 s with a repeat count of 1. The lock mass was enabled for accurate mass measurements. Polydimethylcyclsiloxane (*m/z*, 445.1200025, (Si(CH<sub>3</sub>)<sub>2</sub>O)<sub>6</sub>) ions were used for internal recalibration of the mass spectra.

Raw data files were converted to MGF with Proteome Discoverer software (version 1.2; Thermo Fischer Scientific, San Jose, USA). A precursor mass range of 350 - 5000 Da and signal to noise ratio of 1.5 were the criteria used for generation of peak lists. The peptide and fragment masses obtained were matched automatically against the fungi section of a locally maintained copy of UniprotKB (download 12/2013). MS/MS ion searches were performed using MASCOT Daemon and search engine (version 2.3; Matrix Science, London, UK). The parameters used for database searches included trypsin as a protease with two missed cleavages allowed, and carbamidomethylcysteine, oxidation of methionine, N-terminal protein acetylation and S/T phosphorylation as variable modifications. The tolerance of the ions was set at 5 ppm for parent and 0.8 Da for fragment ion matches. Mascot results were incorporated in Scaffold 3 software (version 3.6, Proteome Software, Portland, USA). Peptide identifications were accepted if they could be established at greater than 95.0% probability as specified by the Peptide Prophet algorithm. Protein identifications were accepted if they could be established at greater than 95.0% probability as specified by the Protein Prophet algorithm.

### 3. Results and Discussion

In asynchronously growing cultures of an SCC1-HA<sub>2</sub> strain treated with the DNA damaging agent methyl methane sulfonate (MMS) or left untreated, no S/T-Q phosphorylation of immunoprecipitated Scc1-HA<sub>2</sub> was detected (**Figure 1(a)** middle panel). Strikingly, in the same experiment, a band migrating between 130 and 170 kD strongly reacted with the anti-P-S/T-Q antibody, only in the presence of MMS (**Figure 1(a)**, top panel). On first analysis, this signal could correspond to either Smc1 or Smc3, the other two major subunits of cohesin, or, alternatively, to two of the three cohesin-associated proteins having a similar size, namely Scc3 or Pds5, the third one, Wpl1, being much smaller (**Figure 1(b)**). Similar observations were made in a *cdc13-1* SCC1-HA<sub>2</sub> strain, in which the temperature-sensitive *cdc13-1* mutation conferred a telomeric damage [19] (**Figure 1(c)**). Moreover, S/T-Q phosphorylation of the same protein, presumably, was also detected in a *cdc13-1* SMC3-Myc<sub>3</sub> strain (**Figure 1(d)**). This indicated that the S/T-Q phosphorylated protein was associated *in vivo* both with Scc1 or Smc3 or was Smc3 itself.

We next set out to perform mass spectrometry analyses (**Figure 2**). To do this, extracts were prepared from asynchronously growing cells of the Scc1-HA<sub>2</sub> strain, submitted to immunoprecipitation with anti-HA antibody and purified proteins separated on an SDS-acrylamide gel and revealed, following transfer onto a nitrocellulose membrane, with the anti-P-S/T-Q antibody, and the anti-HA antibody in parallel (data not shown). The acrylamide gel band corresponding to the P-S/T-Q signal was cut out after superimposing with the reactive nitrocellulose transfer membrane and subjected to trypsin digestion followed by mass spectrometry analysis. In the ana-



**Figure 1.** Scc3 is phosphorylated on S/T-Q residues following DNA damage. (a) Asynchronous cell cultures of an SCC1-HA<sub>2</sub> strain were treated (or not, first lanes) with 0.01% - 0.05% MMS for 2 h at 29°C. Immunoprecipitation-Western blot (IP-W) experiments were performed as indicated; (b) Schematic representation of the cohesin ring and its main associated subunits; (c)-(e) IP-W experiments on *cdc13-1* SCC1-HA<sub>2</sub> cells (c) (34°C is a restrictive temperature for growth), *cdc13-1* SMC3-Myc<sub>3</sub> cells (d) and Myc<sub>2</sub>-PDS5 and Myc<sub>2</sub>-PDS5 SCC1-HA<sub>2</sub> cells (e) as indicated; (f) IP-W experiments on SCC1-HA<sub>2</sub> and SMC1-Myc<sub>18</sub> and SMC3-Myc<sub>9</sub> strains. MW: molecular weight markers; (g) IP-W experiments on SCC1-HA<sub>2</sub> strains with untagged versus 2 Myc-tagged SCC3 as indicated; (h) IP from the SCC1-HA<sub>2</sub> Myc<sub>2</sub>-SCC3 (left undamaged) and Myc<sub>18</sub>-MEC1 (treated with MMS, 2 h to activate its kinase activity symbolized by the asterisk) strains were either mixed together (right lane) or left unmixed (left two lanes as indicated). Arrowhead in top panel indicates the position of the P-S/T-Q reactive band. The SCC1-HA<sub>2</sub> strain also harbored endogenous Myc<sub>2</sub>-SCC3 to verify production of Scc3, the P-S/T-Q reactive protein (bottom panel). Endogenous Myc<sub>18</sub>-Mec1 was also correctly produced (middle panel).



Protein	Number of peptides analyzed	Amino acids coverage	Phosphorylated S/T-Q
Scc3	27	26%	none
Smc1	87	63%	none
Smc3	33	33%	none

(a)

Scc3

MTAVRRSTRI R **TKQGVIEED** YDDEQNTSAQ HVESDKITAK **TSQ**HEEEEDD TGESEESSSE DDYEDQDDDD 0070  
YVDTATAKRR SKRRKPKSAS NTSSKRQKKK PTAQKSAVS HAPAYHRSKK **DQDQYLEIAK** DFQPTFLFDI 0140  
LSTSEDSVIE ELLREWLETY SENRDKFLQE FINLLNCCG SVAR**VEDHDV** HSNESSENETI **GETIQLLFORO** 0210  
**KLHEFYLLIS** KENKRRKNFK **MGPLYQNFAE** FMTKLLEVAN DLQLLYVESD EDD**TSQ**IVTGN LVLLDLTLWS 0280  
SFSVCIRCF RYISTLTLYL PQDYL**TSQ**QAV NLEKNYLAKL SKQLSLEEEK KRPNNK**TLEK** LESTIAE**TSQ** 0350  
**KSVVDSIID** NIVKLCFVHR **YKDVS**DLIRS ESMHLHSIWI KNPYFLKV TFLKYFGWLL SDNSVSVRLQ 0420  
VTK**FLPLII** QNHNSKTDN SAIRQVFERF KTK**ILEVAIR** DVNLDVR**IHS** IQVLTEASSL GYLLDSEILI 0490  
ISSLMFDEEF DPFKTSSFNK RSKFLSTVAK FLARVKEKF DEFIKTHEDL PKEVDGLEVG PVVQVGIFIK 0560  
IINSLDIYHL KDCAEVDSRT KIRML**TSQ**AAE FLSPYISTHL **K**TICNLLISD TESNELIQKL QNSANNNSDD 0630  
EVDDEELDI TPLFPIDR**NS** **TILYLN**VFHG LCAGANNPKI QTKDSVKEIV LPLFYDLLNA ASIESADILC 0700  
PLLESFITFS LDDWISIGYE TELKKITDKT IKAFMDSTIG NSK**VDMKYDI** **FAX**FIIHHH FEKKELQEK**TSQ** 0770  
**LNQIATLK**IH LKKFLQEKMD PNNRSDDYKD LTCSLYELYI NKLTILGRDY PIEVDEELLQ LFLNNFVSR**I** 0840  
**PIIMFQDFDSS** **TAQEIF**NFKML **VLLATW**NLEK WREIEKVRD YENSISKDLR **SVWKPIAAII** GRNLTLVISL 0910  
AATNETFENI NSLFYLKWSA CTSLMDIIVA **IK****IFELK**LPA **DATTWR**YSMS EQFPFYLDHN ASKVLLKIFL 0980  
NYKFLAKQV DVQLERVADE DANLNDLPET GFFENIETEF LLFTVKLGKL **MKLNLDERF** ASRV**ALNKEK** 1050  
**LGPLFKKIVD** DTIMENPEPN **KKNIQ**AKSN **Q****TSQ**REKAPLQ PNSERETDHA NTENNDPDIP MTIDLEPIEE 1120  
**SQ**NNSELAP IEEHPTVDA IDNSDEIT**TSQ**\*

Smc1

MGRVLGLELS NFKSYRGVTK VFGESNFTS IIGPNGSGKS NMMDAISFVL GVRSNHLRSN ILKDLIYRGV 0070  
LNDNSDDYD NEGAASSNPQ SAYVKAFYQK GNKLVELMRI ISRNGDTSYK IDGK**TSYKD** YSIFLENENI 0140  
**LKAKN**FLV FQDVEQIAAQ SPVELSRMFE EVSGSIQYKK EYEELKEKIE KLSKSATESI KNRRRHGEL 0210  
KTYKEGINKN EYRQQLDKK NELQKFQALW QLYHLEQQKE ELTDKLSALN SEISSLKGI NNMKSLQRS 0280  
KSSFVKESAV ISKQSKLDY IFKDKELVS DLRLIKVPOQ AAGKRISHIE KRISLQKDL QROKTYVERF 0350  
**ETQ**LKVVTBS KEAFEEIEKQ SARNYDKFKL NENDLKTYNC LHEKYLTEGG SILEEKIAVL NNDKRIQEE 0420  
LERENKRAID SKRRITEELS **ITGEKLD****TSQ**LDL RVSLNEK NALHTERLHE LKKLSQDIES ANNOEYDLNE 0490  
**KR**ETLVKID **DL**SANORETM KERKLENTIA MLKRFFPGVK GLVHDLCHPK **KEKYGLAVST** ILGNFDSVI 0560  
**VEN**TVAQEC **IAP**LKKORAG TASFIPLDIT ETPLPTLSLP **DQ**QYILSIN AIDYEPEYK AMQYVCGDSI 0630  
ICNTLNIADK LKWKKGIRGK **LV**TIEGALIH KAGLMTGGIS **GDANN**WDKE EYQSLMSLKD **KLIQ**IDELS 0700  
**NGQ**RSNSIRA REVENSVSLL NSDIANLR**TSQ** **V****TSQ**QKSLDE NRLEIKYHND LIEKEIQPKI TELKKLDDL 0770  
ENTKDNLVKE KEALQNNIFK EFTSKIGFTI KEYENHSGEL MRQOSKELQQ LQKQILTVEN KIQFETDRLS 0840  
**TSQ**RYEKAQ KDLNAQVEM KSLEEQEIYAI EMKIGSIESK LEEHKNHLE **LQK**FVTKQS ELNSEDILE 0910  
DMNSNLQVLK RERDGIKEDI EKFDLERVTA LKNCKISNIN IPISSETTID DLPISTDNE AITISNSIDI 0980  
NYKILPKYK ENNTDSARKE LEQK**IE**VEVE ILNELOPNAR ALERYDEAEG RFEVINNETE QLKAEKKIL 1050  
**NOFL**LKIKKR KELFEK**TFDY** VSDHLDAIYR ELTKPNPSNV ELAGGNASLT IEDEDEFFNA GIKYHATPPL 1120  
**KRF**KDMEYLS GGEKTVAALA LLFAINSQOP SPFFVLDEVD AALDITNVQR IAAVIRHRN **PDLO**FIVISL 1190  
**KNT**MFESDSA LVGVYRQQQE **NSSK**ITLIDL SNYAE\*

Smc3

MYIKRVIKG FKTYRNETII DNFSPHQNV IGSNGSGKSN FFAAIRFVLS DDYSNLKREE RQGLIHQGGG 0070  
GVSNSASVEI VFHDPHSMI LPSGVLSSRGD DEVTIRRTVG LKDDYQLND RNVTKGDIVR **MLETAGFSMI** 0140  
**NYE**NIYVQKG IVALTNADK ER**LQ**LEDVV **GAK**SFEVKLK ASLKKMEETE QKKIQINKEM GELNSKLESM 0210  
EQEKKELEKY NELERNRIY QFTLYDRELN EVINQMERLD GDYNNVTYSS EQYIQLODKR EDMIDQVSKK 0280  
LSSTEASLKI KNATDLQAK LRESEI**TSQ**KL TNVNVKIDV QQQIESNEEQ RNLDSATLKE **IKS**IEQRKQ 0350  
KLSKILPRYQ ELTKEEAMYK **LQ**LASLQQKQ **RD**LILKKGEY ARFKSKDERD TWIHSEIEEL KSSIQNLNEL 0420  
**ESQ**LQMDRTS LRQOYSAIDE EIEELIDSIN GPDTKGQLED FDESLIHLKQ KLSSELDTRK ELWRKEQK**LQ** 0490  
**TVLE**TLLSDV **NONQ**RNVNET MSRLANGII NVKEITEKLEK ISPESVFGTL GELIKVNDKY KTCAEVIGGN 0560  
SLFHVVDTE ETATLIMNEL YRMKGGRVTF IPLNRLSLDS DVKFPSNT**TSQ** **Q**IQFTPLIKK IKYEPRFEKA 0630  
VKHVGK**TIV** VKDLGOGLKL AK**HHKL**NAIT LDGDRADKRG VLTGGYLDQH KRTLRLESKN **LNESR****TSQ**HKK 0700  
ILEELDFVRN ELNDIDTRID QVNGNIRKVS NDRESVLTNI EVYH**TS**SLNTR KNEKLILEES LNAIILKLEK 0770  
LNTNRTFAQE KLNTFENDLL QEFDSLSKE EKERLESLETK EISAHNKLN ITSDALEGIT TTIDSLNAEL 0840  
ESKL**IP**END **LES**KMEVGD AFIFGLQDEL KELQLEKESV EKQHENAVLE LGTVQREIES **LIA**ETNNKK 0910  
LLEKANNQOR LLLKKLDNFQ KSVEKTMICK TTLVTRREEL QORIREIGLL PEDALVNDPS DITSQOLLOR 0980  
**LN**DMNTEISG **LK**NVKNKRAE NFKKFNERRK DLAERASELD ESKDSIQDLI VKLKQKQVNA **VDST**FOKVSE 1050  
**NFE**AVFERLV **PR**GAKLIIH RKNNDANDHD SNE**TSQ**KGKDS EIMYTVGSIS VSFNSK**QNEQ** 1120  
**LHVE**QLSGGQ **K**TVCAIALIL AIQMVDPASF YLFDEIDAL DKQY**TSQ**AVAT **LLKEL**SKNAQ FICTTFRTRDM 1190  
LQVADFFRV KYENKISVTI EVNREEAIGF IRGSKNFAEV\*

(b)

**Figure 2.** Mass spectrometry analysis of the anti-P-S/T-Q antibody reactive cohesin band. The acrylamide gel band corresponding to the P-S/T-Q signal was cut out after superimposing with the reactive nitrocellulose transfer membrane of another gel run in parallel and subjected to trypsin digestion followed by mass spectrometry analysis. Smc1, Smc3 and Scc3 peptides were detected in this band. (a) Summary of the results; (b) Amino acid sequences of Scc3, Smc1 and Smc3 showing in yellow the peptides recovered after trypsin digestion (from the same experiment as in (a) and analyzed, and, in red, the amino acids of the S/T-Q type. The coverage of Scc3 sequence following trypsin digestion was rather low, 26%, thus rendering identification of the residues phosphorylated upon DNA damage very difficult.

lyzed sample, three major proteins (among a total of 21) were identified, namely Smc1, Smc3 and Scc3. The number of peptides detected for each of these three proteins, as well the percentage of coverage of the complete sequences, are indicated in **Figure 2(a)** and their sequences shown in **Figure 2(b)**. No event of phosphorylation could be detected in any of these peptides (**Figure 2**). Although Pds5 peptides were not detected, we were nevertheless concerned that it might be implicated in these events. Material immunoprecipitated from a strain bearing endogenous Myc<sub>2</sub>-Pds5 did not bring down the MMS-induced protein with the S/T-Q signal (**Figure 1(e)**). Moreover, no physical interactions between Myc<sub>2</sub>-Pds5 and Scc1-HA<sub>2</sub> either under normal growth conditions or in the presence of 0.05% MMS could be detected (**Figure 1(e)**).

Next, *SMC1* and *SMC3* constructs made in fusion with 9 Myc or 18 Myc epitope tag were integrated at their respective genetic locus, in a strain that also expressed endogenous *SCC1*-HA<sub>2</sub>. The objective was to increase their size so as to see if the P-S/T-Q band was also increased in size (**Figure 1(f)**, top panel). The MMS-induced S/T-Q signal was clearly migrating below the anti-Myc-Smc1-Myc<sub>18</sub> and -Smc3-Myc<sub>9</sub> signals (**Figure 1(f)**, bottom panel) and was of the same size as that in the strains with untagged Smc1 or Smc3 (**Figure 1(a)**). This suggested that the S/T-Q signal was most probably due to Scc3. Confirming this assumption, the S/T-Q band in the Myc<sub>2</sub>-*SCC3* strain was found to migrate above that in the strain with untagged *SCC3* (**Figure 1(g)**), thus definitively establishing that this S/T-Q phosphorylation was indeed on Scc3. We noted that in both strains, the P-S/T-Q reactive material consisted of two bands (**Figure 1(g)**) and closer examination of left panel of **Figure 1(d)**, as well as of **Figure 3(b)**, also suggested the presence of two distinct P-S/T-Q bands. It is therefore possible that two distinct states of S/T-Q phosphorylation on Scc3 exist. However, we do not have at the moment any clue to further document this.

The coverage of Scc3 sequence following trypsin digestion was rather low (**Figure 2(a)**), thus rendering identification of the residues phosphorylated upon DNA damage very difficult. In fact, this mass spectrometric analysis just allowed us to eliminate S14, T348 and T586 (among the 9 possible residues of the S/T-Q consensus sites) as potential candidates for the S/T-Q phosphorylation observed above (**Figure 2(b)**).

It was formally possible that the phosphorylated protein was not Scc3 but rather Pds5 because the established strong binding between the two proteins (see for instance ref. [23]). However, the absence of co-immunoprecipitation of Myc<sub>2</sub>-Pds5 by Scc1-HA<sub>2</sub> and *vice versa* (**Figure 1(e)**) strongly argued that Pds5 was not the Scc1-HA<sub>2</sub>-attached S/T-Q phosphorylated protein.


Next, anti-Myc material IPed from a Myc<sub>18</sub>-*MEC1* strain treated with MMS was mixed *in vitro* with anti-Scc1-HA<sub>2</sub> IP from cells with no incurred damage (**Figure 1(h)**). A phosphorylated S/T-Q band clearly appeared under these conditions, but was absent from either one of the two unmixed immunoprecipitates (**Figure 1(h)**). Although Mec1 and Tel1 are potentially both responsible for phosphorylation on S/T-Q residues, Mec1 rather than Tel1 was chosen for these experiments because genetic inactivation of Mec1 kinase activity alone resulted in total suppression of the P-S/T-Q signal, as shown below.

To further characterize the events described above, we analyzed the S/T-Q response of the *SCC1*-HA<sub>2</sub> strain to other types of damage (**Figure 3(a)**). Hydroxyurea (HU), which activates the replication checkpoint, different from the DNA damage checkpoint, but also dependent on Mec1 kinase, arrests cells in S phase. Nocodazole, a microtubule-poisoning drug, arrests cells in G2/M, like MMS, but in this case cells do not experience DNA damage and the arrest is independent of Mec1. As shown in **Figure 3(b)** (top panel), phosphorylation was detected in immunoprecipitates of Scc1-HA<sub>2</sub> upon treatment with nocodazole, similar to MMS, but not in HU. This suggested that S/T-Q phosphorylation might be related to the G2/M arrest provoked by these drugs rather than to the damage itself. The signal totally disappeared upon genetic inactivation of both *MEC1* and *TEL1* (**Figure 3(b)**, bottom panel), compatible both with the assumption that the phosphorylated residue(s) are indeed the targets of ATR-ATM-like kinases [14] and with the observation that G2/M position in the cell cycle was required for this signal. Both conditions had to be fulfilled as the S/T-Q signal was no longer detected in nocodazole-treated *mec1 tel1* mutant cells even though these cells were still arrested at G2/M (**Figure 3(b)**, bottom). Moreover, in a strain deleted for *MAD2*, in which the spindle checkpoint was no longer functional and which, therefore, no longer arrested at G2/M upon nocodazole-treatment [21], the S/T-Q signal was no longer present (**Figure 3(c)**), thus confirming that damage without concomitant cell cycle arrest in G2/M is not sufficient to trigger phosphorylation of Scc3.


Using again the *SMC1*-Myc<sub>18</sub> and *SMC3*-Myc<sub>9</sub> strains, we could see that the position of the S/T-Q phosphorylated band was at a level different from those for Smc1-Myc<sub>18</sub> and Smc3-Myc<sub>9</sub>, confirming that the detected protein was Scc3 (**Figure 3(d)**, top). Moreover, Myc<sub>2</sub>-Pds5 immunoprecipitate from cells treated with nocoda-

	S	G2 metaphase	metaphase anaphase
MMS 0.05%, 2hr		91%	9%
<i>cdc13-1</i> , 34°C, 2hr		65%	35%
HU 0.4M, 2hr	100%		
Noc. 15mg/ml, 2hr		92%	8%


(a)



(b)



(c)




(d)

**Figure 3.** S/T-Q phosphorylation of Scc3 depends on G2/M arrest and can take place independently of DNA damage. (a) Determination of cell cycle stages upon different treatments by assessing cellular and nuclear morphologies under a fluorescence microscope; (b) Immunoprecipitates from a *SCC1-HA<sub>2</sub>* strain (top) or from a *SCC1-HA<sub>2</sub> mec1 tel1* null strain (bottom), both treated with HU, MMS or nocodazole or left untreated (YPD) were probed with anti P-S/T-Q antibody; (c) Immunoprecipitates from a *SCC1-HA<sub>2</sub>* strain or from a *SCC1-HA<sub>2</sub> mad2* null strain were probed with anti P-S/T-Q antibody; (d) In top panel, strains with the indicated relevant genotypes were treated with nocodazole, processed for *SCC1-HA<sub>2</sub>* IP with anti-HA antibody and probed with anti-P-S/T-Q antibody. Positions of the Myc-tagged proteins relative to the 170 kD marker, not shown here, can be seen in [Figure 1\(f\)](#). In bottom panel, Myc<sub>2</sub>-*PDS5* IP was probed with either anti-Myc or anti-P-S/T-Q antibody following treatment with nocodazole.

zole did not react with the anti-P-S/T-Q antibody ([Figure 3\(d\)](#), bottom).

Next, we assessed the dependence of the DNA damage-induced phosphorylation of Scc3's S/T-Q residue(s) on DNA damage checkpoint genes ([Figure 4\(a\)](#)). First, we could observe that this signal was totally suppressed





**Figure 4.** S/T-Q phosphorylation of Scc3 takes place during mitosis of an unperturbed cell cycle. (a) Schematic representation of the DNA damage checkpoint kinase cascade, highlighting the major components that have been mutated in the present study, as described below (b)-(e); (b) Temperature-sensitive *cdc13-1* mutant cells with the indicated relevant genotype were processed for IP-W experiments as described; (c) Alpha factor-synchronized *SCC1-HA*<sub>2</sub> cells were immunoprecipitated with anti-HA antibody and probed with anti-P-S/T-Q antibody, while the main cell cycle stages were followed in parallel under the fluorescence microscope; (d) (e) Cell cycle analysis of Scc3's S/T-Q phosphorylation following alpha factor block and release in *SCC1-HA*<sub>2</sub> strains harboring the indicated construct (d) or mutations (e); (f) Cell cycle distribution of the strains with the indicated relevant genotype illustrated in panel (e) above. Ordinates indicate the percentages of the different phases of the cell cycle assessed by DAPI staining of nuclei as a function of time following release from a previous alpha factor block in G1.

after expressing a *mec1* kinase-dead mutation from *MEC1* genomic locus (**Figure 4(b)**). Therefore, Mec1 alone appears to be sufficient to phosphorylate Scc3 in response to DNA damage. The signal was also basically suppressed in a *rad24* null background (**Figure 4(b)**), a mutation that prevents loading of the Rad17-Mec3-Ddc1 checkpoint complex and altogether prevents activation of the DNA damage checkpoint kinase cascade. It is worth recalling here that Mec1-Rad53-Chk1 and Rad24-Rad17-Mec3-Ddc1 represent two parallel pathways in the DNA damage checkpoint network and that, although both are essential for checkpoint activation, the Rad24 module is not needed for activation of the Mec1 module. Therefore, in the absence of Rad24, Mec1 was still active, yet the Scc3 S/T-Q signal was absent, thus suggesting that the absence of G2/M arrest in *rad24*Δ cells, rather than the absence of Mec1 activation, was likely responsible for the absence of Scc3 phosphorylation.

Simultaneous inactivation of the *RAD53* and *CHK1* kinases, the two downstream effectors of Mec1 during DNA damage, also resulted in the total disappearance of the Scc3's S/T-Q phosphorylation (**Figure 4(b)**). However, in the single *rad53* mutant, this signal was still present although less intense than in the wild type (**Figure 4(b)**). Activation of Rad53 and Chk1 proceed in parallel and distinct pathways, and activation of the Chk1 pathway has been shown to be particularly evident following *cdc13-1*-induced telomeric damage [24], the type of damage used here (**Figure 4(b)**). Therefore, *cdc13-1 rad53*Δ mutant cells are expected to arrest at least partially because Chk1 is still present, while the *cdc13-1 rad53*Δ *chk1*Δ cells are unable to mark the G2/M arrest, thus explaining the present findings. Since Mec1 is at the top of the DNA damage checkpoint kinase cascade, it was noticeable that the S/T-Q signal, most likely to be caused by the action of Mec1, was absent in the absence of Rad24 or the combined absence of Rad53 and Chk1, two situations in which Mec1 was still present and presumably active (**Figures 4(a)-(b)**). Altogether, these experiments suggested that the S/T-Q signal might be more due to DNA damage-induced cell cycle arrest than to the action of Mec1 itself.

To further document these findings, we next set out to assess Scc3's S/T-Q phosphorylation during a highly synchronous cell cycle (alpha-factor block and release) in the absence of any incurred damage (**Figure 4(c)**). Strikingly, an S/T-Q signal was clearly apparent at 75 - 90 min following release from the G1 block, a time that corresponded to metaphase or the metaphase-anaphase transition based on the simultaneous observation of the sampled cells by microscopy (**Figure 4(c)**). Noticeably, the S/T-Q signal was also present, albeit weaker, before this 75 - 90 min window, as well as after, thus suggesting that S/T-Q phosphorylation of Scc3/cohesin might be important from the onset of mitosis until around telophase (**Figures 4(c)-(e)**). During a synchronous mitosis, the S/T-Q signal was much less intense than that observed in MMS-treated cells or *cdc13-1* cells at 34°C, as expected from the fact that upon one of these two DNA damages the quasi total number of cells in the population eventually arrested at G2/M during the course of the block, 2 h. Again, making use of the *SCC1*-HA<sub>2</sub> strains harboring either endogenous *SMC1*-Myc<sub>18</sub> or *SMC3*-Myc<sub>9</sub>, established that, by elimination, Scc3 was most probably the protein phosphorylated on S/T-Q during mitosis of undamaged synchronized cells (**Figure 4(d)**). Finally, we could demonstrate that the cell cycle-associated S/T-Q phosphorylation of Scc3/cohesin was dependent on the presence of Mec1, but that simultaneous genetic inactivation of *RAD53* and *CHK1* did not prevent its occurrence (**Figures 4(e)-(f)**).

To our knowledge, in *S. cerevisiae*, only Scc3's S343 has been reported to be phosphorylated upon DNA damage (ref. [25] and see also <http://www.phosphogrid.org>). The potential significance of this phosphorylation event has not been established. Other phosphorylation events on residues of *S. cerevisiae* Scc3, namely S28, S34 and S623, have been reported, but these were labeled as ambiguous; phosphorylation on Scc3's S628 has also been found (<http://www.sbeams.org>). In human cells, Plk1 phosphorylates SCC3/SA2, a step essential in the cohesin dissociation pathway during prophase and prometaphase [26]. In budding yeast, the major response of cohesin to DNA damage is on its Scc1/Mcd1 subunit [11], while in humans, in case of DNA damage, phosphorylation occurs on Smc1 and Smc3, and is thought to be mediated directly by the kinases ATM and ATR [12] [13]. Scc3 has major roles in cohesin-chromosomes association/dissociation, as well as in maintaining sister chromatid cohesion, in cooperation with the other regulatory cohesin subunits Pds5 and Wapl [5] [23] [27].

In summary, we have uncovered here an event of phosphorylation of cohesin on one or several S/T-Q residues (potentially by the *MEC1*/ATR kinase) that most probably takes place on Scc3, a protein closely associated with the cohesin ring. Most importantly, this event takes place during mitosis of an unperturbed cell cycle, culminating in intensity around anaphase and telophase, but also present a little earlier and later. This event is of prime importance for two reasons. First, given the key position of Scc3 at the heart of the cohesin complex, this finding suggests that Scc3 phosphorylation might represent a key element in coupling the assembly/disassembly reactions of the Smc1-Smc3-Scc1 cohesin ring with the other cohesin regulatory subunits as well as with mitotic

events. Second, the present findings suggest that Mec1 might intervene during mitosis independently of any incurred DNA damage and therefore independently of its checkpoint function to control cohesin through direct regulation of Scc3 by phosphorylation on S/T-Q residues. Presumably, the P-S/T-Q signal detected upon treatment with MMS or nocodazole corresponds to a G2/M enrichment of the cell population as it disappears when the damaged-induced cell cycle arrest is compromised by a mutation in a checkpoint gene. Further analyzing this Scc3 phosphorylation is crucial and represents a future challenge to fully understand cohesin biology.

## Acknowledgements

We thank Akira Matsuura, Andrew Emili, Leland Hartwell, Wolf-Dietrich Heyer, Andrew Murray and Errol Friedberg for the gift of strains and plasmids. We thank Valérie Labas and Grégoire Harrichaux for performing the mass spectrometry analysis at the “Plate-forme d’Analyse des Biomarqueurs” (INRA, Nouzilly, France). The high resolution mass spectrometer was financed (SMHART project) by the European Regional Development Fund (ERDF), the “Conseil Régional du Centre”, the French National Institute for Agricultural Research (INRA) and the French National Institute of Health and Medical Research (Inserm). This work was financially supported by subventions from the “Centre National de la Recherche Scientifique” and the “Université François-Rabelais Tours” to the UMR CNRS 7292/University of Tours-GICC Research Unit.

## References

- [1] Guacci, V., Koshland, D. and Strunnikov, A. (1997) A Direct Link between Sister Chromatid Cohesion and Chromosome Condensation Revealed through the Analysis of *MCD1* in *S. cerevisiae*. *Cell*, **91**, 47-57. [http://dx.doi.org/10.1016/S0092-8674\(01\)80008-8](http://dx.doi.org/10.1016/S0092-8674(01)80008-8)
- [2] Michaelis, C., Ciosk, R. and Nasmyth, K. (1997) Cohesins: Chromosomal Proteins That Prevent Premature Separation of Sister Chromatids. *Cell*, **91**, 35-45. [http://dx.doi.org/10.1016/S0092-8674\(01\)80007-6](http://dx.doi.org/10.1016/S0092-8674(01)80007-6)
- [3] Ciosk, R., Shirayama, M., Shevchenko, A., Tanaka, T., Toth, A., Shevchenko, A. and Nasmyth, K. (2000) Cohesin's Binding to Chromosomes Depends on a Separate Complex Consisting of Scc2 and Scc4 Proteins. *Molecular Cell*, **5**, 1-20. [http://dx.doi.org/10.1016/S1097-2765\(00\)80420-7](http://dx.doi.org/10.1016/S1097-2765(00)80420-7)
- [4] Uhlmann, F., Wernic, D., Poupert, M.A., Koonin, E.V. and Nasmyth, K. (2000) Cleavage of Cohesin by the CD Clan Protease Separin Triggers Anaphase in Yeast. *Cell*, **103**, 375-386. [http://dx.doi.org/10.1016/S0092-8674\(00\)00130-6](http://dx.doi.org/10.1016/S0092-8674(00)00130-6)
- [5] Peters, J.M., Tedeschi, A. and Schmitz, J. (2008) The Cohesin Complex and Its Roles in Chromosome Biology. *Genes and Development*, **22**, 3089-3114. <http://dx.doi.org/10.1101/gad.1724308>
- [6] Nasmyth, K. (2011) Cohesin: A Catenase with Separate Entry and Exit Gates? *Nature Cell Biology*, **13**, 1170-1177. <http://dx.doi.org/10.1038/ncb2349>
- [7] Ocampo-Hafalla, M.T. and Uhlmann, F. (2011) Cohesin Loading and Sliding. *The Journal of Cell Science*, **124**, 685-691. <http://dx.doi.org/10.1242/jcs.073866>
- [8] Simmons-Kovacs, L.A. and Haase, S.B. (2010) Cohesin: It's Not Just for Chromosomes Anymore. *Cell Cycle*, **9**, 1750-1753. <http://dx.doi.org/10.4161/cc.9.9.11792>
- [9] Losada, A. (2014) Cohesin in Cancer: Chromosome Segregation and Beyond. *Nature Reviews Cancer*, **14**, 389-393. <http://dx.doi.org/10.1038/nrc3743>
- [10] Mehta, G.D., Kumar, R., Srivastava, S. and Ghosh, S.K. (2013) Cohesin: Functions beyond Sister Chromatid Cohesion. *FEBS Letters*, **587**, 2299-2312. <http://dx.doi.org/10.1016/j.febslet.2013.06.035>
- [11] Heidinger-Pauli, J.M., Unal, E., Guacci, V. and Koshland, D. (2008) The Kleisin Subunit of Cohesin Dictates Damage-Induced Cohesion. *Molecular Cell*, **31**, 47-56. <http://dx.doi.org/10.1016/j.molcel.2008.06.005>
- [12] Kim, S.T., Xu, B. and Kastan, M.B. (2002) Involvement of the Cohesin Protein, Smc1, in Atm-Dependent and Independent Responses to DNA Damage. *Genes and Development*, **16**, 560-570. <http://dx.doi.org/10.1101/gad.970602>
- [13] Yazdi, P.T., Wang, Y., Zhao, S., Patel, N., Lee, E.Y. and Qin, J. (2002) SMC1 Is a Downstream Effector in the ATM/NBS1 Branch of the Human S-Phase Checkpoint. *Genes and Development*, **16**, 571-582. <http://dx.doi.org/10.1101/gad.970702>
- [14] Kim, S.T., Lim, D.S., Canman, C.E. and Kastan, M.B. (1999) Substrate Specificities and Identification of Putative Substrates of ATM Kinase Family Members. *The Journal of Biological Chemistry*, **274**, 37538-37543. <http://dx.doi.org/10.1074/jbc.274.53.37538>
- [15] Grandin, N. and Reed, S.I. (1993) Differential Function and Expression of *Saccharomyces cerevisiae* B-Type Cyclins in Mitosis and Meiosis. *Molecular and Cellular Biology*, **13**, 2113-2125.

- [16] Emili, A. (1998) MEC1-Dependent Phosphorylation of Rad9p in Response to DNA Damage. *Molecular Cell*, **2**, 183-189. [http://dx.doi.org/10.1016/S1097-2765\(00\)80128-8](http://dx.doi.org/10.1016/S1097-2765(00)80128-8)
- [17] Siede, W., Nusspaumer, G., Portillo, V., Rodriguez, R. and Friedberg, E.C. (1996) Cloning and Characterization of RAD17, a Gene Controlling Cell Cycle Responses to DNA Damage in *Saccharomyces cerevisiae*. *Nucleic Acids Research*, **24**, 1669-1675. <http://dx.doi.org/10.1093/nar/24.9.1669>
- [18] Bashkirov, V.I., Bashkirova, E.V., Haghazari, E. and Heyer, W.D. (2003) Direct Kinase-to-Kinase Signaling Mediated by the FHA Phosphoprotein Recognition Domain of the Dun1 DNA Damage Checkpoint Kinase. *Molecular and Cellular Biology*, **23**, 1441-1452. <http://dx.doi.org/10.1128/MCB.23.4.1441-1452.2003>
- [19] Garvik, B., Carson, M. and Hartwell, L. (1995) Single-Stranded DNA Arising at Telomeres in *cdc13* Mutants May Constitute a Specific Signal for the RAD9 Checkpoint. *Molecular and Cellular Biology*, **15**, 6128-6138.
- [20] Takata, H., Kanoh, Y., Gunge, N., Shirahige, K. and Matsuura, A. (2004) Reciprocal Association of the Budding Yeast ATM-Related Proteins Tel1 and Mec1 with Telomeres *In Vivo*. *Molecular Cell*, **14**, 515-522. [http://dx.doi.org/10.1016/S1097-2765\(04\)00262-X](http://dx.doi.org/10.1016/S1097-2765(04)00262-X)
- [21] Li, R. and Murray, A.W. (1991) Feedback Control of Mitosis in Budding Yeast. *Cell*, **66**, 519-531. [http://dx.doi.org/10.1016/0092-8674\(81\)90015-5](http://dx.doi.org/10.1016/0092-8674(81)90015-5)
- [22] Grandin, N., Corset, L. and Charbonneau, M. (2012) Genetic and Physical Interactions between Tel2 and the Med15 Subunit of Mediator in *Saccharomyces cerevisiae*. *PLoS ONE*, **7**, e30451. <http://dx.doi.org/10.1371/journal.pone.0030451>
- [23] Kulemzina, I., Schumacher, M.R., Verma, V., Reiter, J., Metzler, J., Failla, A.V., Lanz, C., Sreedharan, V.T., Rätsch, G. and Ivanov, D. (2012) Cohesin Rings Devoid of Scc3 and Pds5 Maintain Their Stable Association with the DNA. *PLoS Genetics*, **8**, e1002856. <http://dx.doi.org/10.1371/journal.pgen.1002856>
- [24] Sanchez, Y., Bachant, J., Wang, H., Hu, F., Liu, D., Tetzlaff, M. and Elledge, S.J. (1999) Control of the DNA Damage Checkpoint by Chk1 and Rad53 Protein Kinases through Distinct Mechanisms. *Science*, **286**, 1166-1171. <http://dx.doi.org/10.1126/science.286.5442.1166>
- [25] Chen, S., Albuquerque, C.P., Liang, J., Suhandynata, R.T. and Zhou, H. (2010) A Proteome-Wide Analysis of Kinase-Substrate Network in the DNA Damage Response. *The Journal of Biological Chemistry*, **285**, 12803-12812. <http://dx.doi.org/10.1074/jbc.M110.106989>
- [26] Hauf, S., Roitinger, E., Koch, B., Dittrich, C.M., Mechtler, K. and Peters, J.M. (2005) Dissociation of Cohesin from Chromosome Arms and Loss of Arm Cohesion during Early Mitosis Depends on Phosphorylation of SA2. *PLoS Biology*, **3**, e69. <http://dx.doi.org/10.1371/journal.pbio.0030069>
- [27] Roig, M.B., Löwe, J., Chan, K.L., Beckouët, F., Metson, J. and Nasmyth, K. (2014) Structure and Function of Cohesin's Scc3/SA Regulatory Subunit. *FEBS Letters*, **588**, 3692-3702. <http://dx.doi.org/10.1016/j.febslet.2014.08.015>